



Adam J. Farber, MD

Sports Medicine and Orthopaedic Surgery
Board Certified; Fellowship-trained in Sports Medicine & Arthroscopic Surgery
P: 480-219-3342; F: 480-219-3271

POST-OPERATIVE INSTRUCTIONS CLAVICLE FRACTURE FIXATION

Wound Care

The original post-operative dressings generally should be left in place for 7 days. Do not remove the steri-strips or cut any of the visible sutures. Cover the incision with clean dry gauze. Keep the incision clean and dry. Bruising down into the elbow and chest wall is not uncommon.

Be sure to watch for signs and symptoms of infection after surgery, which includes: redness, increased swelling, warmth, wound drainage, or a fever greater than 101.5°. If you notice any of these signs or symptoms, or if you are experiencing discoloration of the fingers or numbness/tingling, please notify Dr. Farber's office.

Showering and Dressing

Showering is allowed with plastic covering the wounds. You may shower by placing a large garbage bag over your sling starting the day after surgery. DO NOT immerse your incision under water. This means NO baths, swimming, or hot tubs of any kind for at least two weeks after surgery. To clothe yourself, remember to put your operative arm in your shirt first (keeping your hand at your side), pull the shirt in position and then reach through the remaining sleeve with your good arm. A helpful hint: button up shirts are the easiest and safest to wear while recovering from surgery. Once dressed, be sure to properly place your operative arm in the shoulder sling.

Sleeping

Patients are generally more comfortable sleeping in a reclining chair or with pillows propped behind the shoulder and/or under the forearm. Some difficulty with sleeping is common for several weeks after surgery.

Ice

Icing is very important for the first 5-7 days after surgery. Ice should be applied for 20-minute periods 3-6 times per day. Care must be taken with icing to avoid frostbite to the skin. To avoid frostbite, place a towel or T-shirt between the ice and your skin.

Sling

Your sling should be worn essentially at ALL times including sleeping (except for hygiene and exercises). Remove the sling 3 times per day to work on gentle elbow range of motion exercises. Minimize shoulder range of motion until your first post-operative follow-up appointment. Work on wrist, hand, and finger range of motion while wearing your sling.

Activity

Be sure to use and move your hand, wrist, and elbow in order to decrease swelling in your arm. Squeezing a tennis ball or squeeze ball is encouraged as it will reduce the amount of swelling in your arm and hand. While exercise is important, don't over-do it.

Exercises

- **Range of Motion Exercises:** While maintaining your arm at your side, begin elbow, wrist, and hand range of motion exercises within 24 hours after surgery. Perform these exercises 3 times per day.

Medications

Take as prescribed. **Narcotic pain medications** : Percocet is used for severe pain. It can be taken up to every four hours as necessary. Most patients only require Percocet for the first week. Once pain is better controlled, you may simply take extra strength Tylenol one to two tabs every six hours. Take these medications with food. If you have any problems taking the medications please stop them immediately and notify the office.

Local anesthetics are put adjacent to the incision during surgery. It is not uncommon for patients to encounter more pain on the first or second day after surgery when the effect of these medications wears off. Using the pain medication as directed will help control pain with little risk of complication. Taking pain medication before bedtime will assist in sleeping. It is important not to drink while taking narcotic medication. You can supplement the narcotic medications with 200 mg or 400 mg of ibuprofen every 4-6 hours. You should resume your normal medications for other conditions the day after surgery. I have no specific diet restrictions after surgery but extensive use of narcotics can lead to constipation. High fiber diet, lots of fluids, and muscle activity can prevent this occurrence.

Eating

The anesthetic drugs used during your surgery may cause nausea for the first 24 hours. If nausea is encountered, drink only clear liquids (i.e. Sprite or 7-up). The only solids should be dry crackers or toast. If nausea and vomiting become severe or the patient shows signs of dehydration (lack of urination) please call the doctor or the surgical center.

Driving & Work

NO driving is allowed while on narcotic medications. In addition, the operative arm should not be used on the steering wheel until allowed by your physician. You may return to sedentary work ONLY or school 4-5 days after surgery, if pain is tolerable.

Follow-Up

Your initial follow up visit will usually be 10-14 days after surgery. If you have any questions, concerns or problems please feel free to contact the office at (480) 219-3342.